

Small Planted Tanks for Pet Shrimp

by Diana Walstad
(updated May 2023)

Setting up a planted fish tank is littered with pitfalls. Newly purchased plants have to adapt to a new substrate, light source, and water conditions. Emergent-grown plants will have to adapt to the submerged condition. Chances are some plant species will not survive. Algae may become a problem. Fish add another layer of complication. Too often purchased fish get diseases that prompt unwary hobbyists to add chemicals that injure plants.

Small bowls for shrimp are much less prone to problems and frustration. In this article, I describe two methods for setting up small planted tanks for pet shrimp. The Bowl Setup is quick and easy. The Dry Start Method (DSM) is more complicated, but it has some major advantages over the usual (submerged) startup.

It was only when I started keeping shrimp as pets that I saw the advantage of scaling down to bowls and tiny tanks. I wish I had done so earlier. The shrimp—Red Cherry Shrimp (or RCS)—are cute, inexpensive and low maintenance (**Fig 1**). They are perfect for small planted tanks and bowls—no heater, no filter, no special foods, and no fish diseases. Water changes are easy, because you're only working with one or two gallons.

Small shrimp tanks are a great way to start out with aquarium plants. Beginners learn how to work with soil. They discover which plant species adapt best to their aquarium conditions before setting up a large tank.

Learning how to grow aquarium plants is worth the effort. Plants purify the water and substrate, thereby reducing tank maintenance (water changes, gravel vacuuming, etc).¹

Bowl Setup Procedure

Materials:

- Glass bowls [1 gal (3.8 liters), sold by WalMart as 'fish bowls' and made by Anchor Hocking]
- Soil- Potting soil used for growing houseplants (**Fig 2**)
- Sand- pool-filter sand or "play sand" from hardware stores
- Plants- I chose small plants that for me are reliable growers (**Fig 3**)



Fig 1. Red Cherry Shrimp (RCS) or *Neocaridina davidi*

This adult female, which is about 3/4" in length, is brightly colored. Adult males are smaller, slimmer, and usually lighter-colored. RCS are less demanding than other pet shrimp.

¹ My book *Ecology of the Planted Aquarium* (4th Ed, 2023) explains how plants make aquarium keeping easier. For example, Chapter II ('Plants as Water Purifiers') discusses plant uptake of heavy metals, ammonia, and nitrite. Chapter III ('Bacteria') discusses toxin processing by soil bacteria. Chapter VI ('Plant Nutrition and Ecology') documents the preferential uptake of toxic ammonia by aquatic plants.

- Water- I treated it with an aquarium water conditioner that neutralizes heavy metals. (Shrimp are quite sensitive to heavy metals, more so than fish. The 0.8 ppm zinc in my well water will kill shrimp.)
- Light- window light and a 10.5” Clamp Light with a screw-in 14 watt CFL (compact fluorescent light)
- Day length was 14 hr (hours) with a 2-4 hr afternoon Siesta of lower light beginning at noon.²
- Temperature- I set these bowls up in the summer when temperatures (72- 82°F) were ideal. During the winter, I insert a heat source (i.e., Hydor Mini-heater, a 7.5 watt reptile heating pad). RCS can survive low temperatures (~55°F), but plants won’t grow.

Procedure:

I first remove sticks and larger wood pieces from the bagged potting mix (**Fig 2**). Then, I place 2 cups of it into the bottom of the bowl such that the depth is no more than 1 inch. I then add about 2/3 cup water—enough to moisten the soil but not make it soupy. I let the soil sit overnight so that the the soil’s air pockets fill with water, thereby making soil particles less likely to float and planting easier.

The next day as I put the plants into the soaked soil, I cover the soil and plant roots with ~1 cup sand. I use just enough sand to hold down the soil particles. (A thin covering will be less likely to block oxygen entry into the soil from the overlying water. For if the soil becomes too anaerobic, it will kill plant roots.)

I add water carefully so as not to disturb the soil. I always use my hand (or some other object) to block the main force of the incoming water. Then, I make minor adjustments such as pulling out leaves buried by the sand, adding more stem plants, and spooning more sand over areas with escaping soil. Sometimes, I add a small stone to hold a plant down temporarily. The first water usually has some floating soil particles, so I change water once or twice until it is reasonably clear. (Minor turbidity should go away by the next day leaving the water crystal clear.)

Sometimes, I add shrimp right away, but I advise waiting a couple weeks. That’s because the submergence of any terrestrial soil induces chemical and biological chaos within the soil (my book, Ch VII). During the first few weeks, there is a temporary but explosive release of nutrients—including NH₃ (ammonia). And as soil bacteria feed frenetically on all that ‘fresh’ organic matter, they pull oxygen out of the water. Plant growth won’t be sufficient yet to counter these problems. Thus, hobbyists adding shrimp immediately after setup need to closely monitor ammonia levels, prepare for water changes and possibly add aeration for the first couple weeks.



Fig 2. Potting Mix

Almost any potting soil will work. This particular potting mix contains well-composted organic matter and has a low NPK ratio of 0.10:0.05:0.05. Soil containing excessive chemical fertilizers like nitrates and sulfates will generate toxins (e.g., nitrite and hydrogen sulfide) upon submergence.

² Plants need a day length of at least 12 hr to mimic summer growing conditions. Shorter day lengths signal the onset of winter and trigger slower plant growth [1]. A siesta regenerates CO₂ levels so that plants have some CO₂ for afternoon photosynthesis (my book, Fig X-1, p. 192). It balances light and CO₂ for better plant growth.

Bowls have been without problems (**Figs 3 and 4**). Plants grew well ‘right-off-the-bat’.

Fig 3. Newly Setup Shrimp Bowl

I used plants (surplus from other tanks) that I knew would do well. Rosette and grass-like plants included *Sagittaria subulata*, *S. graminea*, *Echinodorus tenellus*, and *E. radicans* (dwarf). For stem plants, I used *Bacopa monnieri* and *Rotala rotundifolia*. I later threw in some *Ludwigia arcuata*, Java moss (*Vesicularia dubyana*) and *Riccia fluitans*. For these first ‘beginner bowls’, I nixed *Cryptocoryne* species, which can melt down during adjustment to a new environment. I did not add ‘furniture’ (e.g., driftwood) that would smother the soil layer, decrease the planted area, and possibly leach nutrients into the water.



Fig 4. Established Bowls

Photo shows the two bowls at 7 months (Jan 2010). Plant growth was rapid from the beginning so that the bowls positively sailed through the startup period. *Bacopa monnieri* is blooming and growing emergent outside one bowl. I’ve had to do very little maintenance except minor plant pruning and water top-offs. The bowl on the right has a little mat alga that I pull out with tweezers.



DSM Setup Procedure

In 2009, I set up a couple tanks for shrimp with tiny, ground-covering “carpet plants” using the DSM (Dry Start Method) [2]. Startup involves growing plants emergent under terrarium-like conditions (**Fig 5**). Only after the plants became firmly established and spreading (usually takes about 6 weeks), did I submerge them fully (I added water gradually to the tanks over a period of several days.).

The DSM has several advantages over the usual setup method. Emergent-grown plants often grow 4-10 faster than plants grown submersed. Plants quickly develop an extensive root system. Because plants are not submerged, they don’t have to compete with algae. Moreover, emergent-grown plants

don't have to adapt to the submerged condition and grow all new leaves. Meanwhile, the soil will have gone through several weeks of decomposition (or "mineralization") before it is submerged; it will be more stable and less likely to release large quantities of algae-stimulating nutrients into the water.

For the DSM tanks, I worked with some of the obstacles that handicap beginners. That is, I started with purchased plants that were in their emergent form. In addition, I used either unfamiliar plant species or those that had not done well for me in the past in my larger fish tanks.

Because of the emergent startup, I was less cautious about the soil layer. For example, I mixed soil types, by combining a mineral soil (i.e., yard dirt) with an organic soil (i.e., potting soil). Under these conditions, the yard dirt would release its copious iron into the water and greatly stimulate algae. However, algae is not an issue for a DSM startup. Once I submerged the tank, I planned to include floating plants, which would need iron in the water, and ordinary mineral soils are full of it.

Emergent plants actually grow better in a mineral soil than an organic soil [3]. However, I needed soil organic matter to provide CO₂ to the plants after submergence. (The decomposition of soil organic matter releases copious CO₂ into the water.) I also added a *little* fertilizer to the soil layer.

Materials and Key Factors:

- Two-gal tanks
- Glass lids to keep in warmth and moisture
- Mineral soil (garden soil); I used the southeastern Piedmont clay topsoil from my property.
- Organic potting soil (i.e., Miracle Gro's Organic Choice 'Potting Mix' shown in Fig 2)
- Plant tabs containing *small* amounts of KNO₃, NH₄H₂PO₄, etc in a clay filler
- Sand- pool filter sand or "play sand" from a hardware store
- Water- tapwater treated with an aquarium water conditioner to neutralize heavy metals
- Light- 10.5" Clamp Light with a 14 watt screw-in CFL. The clamp light rests directly on the glass lids of both tanks.



Fig 5. DSM (Dry Start Method) Tanks at Startup

I took this overhead photo of the two 2-gal tanks soon after setup. Plants are identified in the photo with red lettering and the following abbreviations: AN = *Anubias nana* (petite), HC = *Hemianthus callitrichoides*; EA = *Eleocharis acicularis*; and GE = *Glossostigma elatinoides*. Tank A is at the top.

- Day length- During the emergent phase, I kept lights on continuously for 14 hr per day.³ After submergence, I put tanks on the “Siesta Regimen” that I used for the bowls.
- Heaters (after submergence)- 7.5 watt Hydor Mini-heaters (reptile heating pads)
- Air bubblers (after submergence) that released a single air bubble at a time to provide *gentle* water circulation without driving off precious CO₂
- Plants- Most were “carpet plants” such as *Anubias nana* (petite); *Hemianthus callitrichoides*; Dwarf Hairgrass (*Eleocharis acicularis*); *Glossostigma elatinoides*; and Four Leaf Clover (*Marsilea quadrifolia*). I ordered these tiny, ground-covering species on-line from AquariumPlants.com. (Plants, which arrived in excellent condition, were originally grown by Florida Aquatic Plant Nurseries.) Plants were in their emergent form. After submergence, I added some of my own surplus plants [floating Frogbit (*Limnobium laevigatum*); *Hemianthus micranthemoides*; and *Riccia fluitans*].

Procedure:

Because I would be planting small delicate plants, I used a shallow, fine-textured substrate. I first removed sticks and larger wood pieces from the bagged potting mix. To Tank A, I added 4 cups garden soil and then 1.5 cups potting mix. To Tank B, I added 2 cups garden soil and 3.5 cups potting mix. I did not mix the soils and made sure the potting mix went on top where it would get more oxygen. I crushed and sprinkled two plant tab fertilizers over the soil in each tank.

I then added 1 to 2 cups of aged aquarium water—enough to thoroughly moisten the soil but not make it soupy. I drained off any excess. I spooned in some washed sand and planted the plants.

For the next 10 weeks, I kept the glass lids snug and made sure that the soil stayed moist. Every week or so, I added a little water. Most mornings, the tanks had condensed water on the glass showing that the tanks were sufficiently humidified. Towards the end when plants were growing rapidly but turning yellow, I added a couple crushed plant fertilizer tabs to the replenishment water.



Fig 6. DSM Tanks at 10 Weeks

HC and EA have spread considerably since the initial planting. However, within a few days of planting, the GE turned brown and died. The leaves of *Marsilea quadrifolia* turned brown (as shown here in the bottom tank). The AN grew very slowly. Photo was taken just before I submerged plants. ‘Tank A’ is at the top; ‘Tank B’ at the bottom. See Fig 5 for plant abbreviations.

³ For emergent plants, the more light, the better. They can use light intensity up to full sunlight for their photosynthesis. In contrast, submerged plants can use only about one-tenth of full sunlight (my book, Fig VIII-2, p. 150).

At 10 weeks, I finally added water and submerged the plants. I changed 100% of the water twice within the next two days to wash out leftover fertilizers.

A week later, I added RCS to Tank A and Grass Shrimp to Tank B. I probably could have added them sooner, but I wasn't sure what would happen.

Tank Results:

During the 10 weeks prior to submergence, the *H. callitrichoides* and *E. acicularis* multiplied well and formed an extensive root system. Submergence did not seem to hurt them, and they did not shed their leaves. Indeed, they seemed to thrive after submergence (**Figs 7 and 8**). *M. quadrifolia* actually seemed to grow better with less leaf-browning after I submerged it.

Tanks dominated by small delicate plants, are extremely vulnerable to algae. Within ten days of submergence, I noticed some alga growth on the glass sides. I took quick action: (1) cleaned the glass; (2) changed water; (3) raised the Clamp Light a few inches; (4) added snails; and (5) added floating Frogbit and *Riccia fluitans*. After the Frogbit started growing well (**Fig 8**) and the algae retreated, I lowered the Clamp Light back down to where it had been before—resting on the glass lids. Once the Frogbit began to grow, the algae retreated.⁴

Tank A with 27% Potting Mix has less organic matter in the substrate than Tank B with 64% Potting Mix. The difference didn't seem to matter.

Discussion

The shrimp bowls were unheated. I was concerned that the plants and my RCS might do poorly during the winter. Night-time room temperatures were between 60 and 65°F, and



Fig 7. DSM Tanks after Submergence

Three weeks after submergence, carpet plants are growing very well. They have made such a thick mat, I can pour water into the tank without disturbing the soil layer. Tank on the right has RCS, while the tank on the left contains native Grass Shrimp.



Fig 8. Frogbit to the Rescue!

I removed the tank lids to get this photo (at 6 weeks post-submergence) showing the Frogbit floating on the water surface (you can also see floating mats of *Riccia*). Algae threatened to take over the tanks within days of submergence. I consider Frogbit vital for controlling algae in these two tanks.

⁴ Emergent plants and floating plants have major advantages over algae that the hobbyist can exploit (my book, Ch VII, the 'Aerial Advantage').

daytime temperature rarely got above 70°F. However, RCS in their native habitat are exposed to temperatures between 6°C (43°F) and 30°C (86°F) [4].

So my concern was more for the plants than the shrimp. The optimum temperature range for most aquarium plants is 72- 82°F, but the lower winter temperatures did not seem to affect plant growth in the bowls. Many species were temperate plants (e.g., *Bacopa monnieri*) that—unlike plants from the tropics—tolerate lower temperatures. When in January, I measured day-time CO₂ levels in the bowls, I found rapid daytime CO₂ depletion comparable to what I measured in September. This CO₂ depletion means that some plants—despite the cooler temperatures—were actively photosynthesizing.

As for the 2-gal tanks.... They were coddled. I heated them with a mini-heater and used *very* gentle air bubbling to help circulate the heater's warmth but not drive off CO₂. The temperature did not go below 70°F in these two tanks. With the onset of winter, RCS in the heated tank were the only ones to have babies. And I routinely saw baby shrimp attached to the mini-heater. {Scientists investigating RCS generally maintain their colonies at 27°C (80.6°F) [5]}.

For a couple weeks after submergence, I ran Tank B without a heater. The carpet plants looked fine. However, I noticed that the Frogbit was yellow and not multiplying, whereas in Tank A with a heater, it was thriving. Since these tanks depend on Frogbit to control algae, I quickly added a heater and air-bubbler, identical to what I have in Tank A. Frogbit turned green and started multiplying. [This Frogbit was a tropical species (*Limnobium laevigatum*), not the native Frogbit (*Limnobium spongia*) common to the Southeastern U.S.A. that grows well at lower temperatures.]

Beginners often start with unfamiliar and newly purchased plants. Most vendors sell plants in their emergent form. Some plant species adjust to submergence better than others, but they still have to adjust. If plants are not growing well by the first couple of weeks and algae becomes entrenched, the tank may not succeed. That's because a large alga mass will quickly remove all CO₂ from the water,⁵ making it very difficult for the plants.

I set up the DSM tanks to see how well the method worked with some challenging plants, that is, "carpet plants." I consider them difficult for a low-tech setup where there's no artificial CO₂ injection.⁶ These plants have not done well for me in large fish tanks where they had to compete with *Cryptocoryne* and *Echinodorus* species. However, in small tanks by themselves and under conditions that minimize CO₂ loss, they did surprisingly well.

Carpet plants *H. callitrichoides* and *E. acicularis* started via the DSM did fine. Moreover, the emergent startup period required almost no maintenance (or worry). For 10 weeks all I did was occasionally add a little water to moisten the soil and keep the air humidified.

G. elatinoides and *M. quadrifolia* had problems during the DSM startup (**Fig 6**). I suspect that ethylene gas, a plant hormone produced in large amounts by wounded and/or stressed plants, caused those problems. Ethylene can cause leaf-browning and death, thereby inducing even more ethylene release. The two affected plant species may have been more sensitive to ethylene, or they were more damaged during planting than the other plant species. Air circulation helps dilute the released gas. Should I try DSM again, I will handle the plants more gently, remove dying leaves immediately, and air out the tanks more during the emergent phase.

I do not recommend carpet plants for beginning hobbyists. The plants are not that competitive. I started my carpet plants under ideal (i.e., emergent) conditions and then provided them with a

⁵ Algae can grow at lower CO₂ levels [6] and use bicarbonates more effectively than plants [7].

⁶ High-tech tanks have CO₂ injection. There's enough CO₂ for all plants, including those that are less-competitive in obtaining CO₂. In my tanks, plants compete for a limited supply, so not every plant species is going to do well.

carefully controlled tank. For example, the larger, more robust “baby tears” *Hemianthus micranthemoides* in one tank eventually crowded out the delicate carpet plant *H. callitrichoides*. Carpet plants will have a better chance if they are accompanied by floating plants. Carpet plants—on their own—cannot remove nutrients sufficiently from the water to prevent algae (or purify the water for the shrimp). Floating plants protect carpet plants without competing with them for CO₂. (Floating plants get their CO₂ from the air, not the water, so—unlike submerged plants—they are not CO₂-limited.)

My DSM tanks required more maintenance than the bowls. I had to change water at least once every week during the first 6 weeks following submergence. Occasionally, I had to remove small algae mats (using a toothbrush) that threatened to spread over the plant carpet. After the Frogbit and *Riccia* started growing, the tanks stabilized and did quite well with minimal maintenance.

For the more advanced hobbyist who wants to experiment, the DSM is an interesting option. If it works for finicky carpet plants, it should work well for other aquarium plants. Emergent growth is common for aquatic plants during the dry season in their native habitats. The vast majority of aquarium plants (species of *Anubias*, *Echinodorus*, *Sagittaria*, *Cryptocoryne*, *Bacopa*, *Ludwigia*, *Rotala*, *Myriophyllum*, *Microsorium*, etc) can be grown emergent, and therefore, lend themselves to a DSM startup. Only a few aquatic plants do not have an emergent form; these include species of *Aponogenton*, *Najas*, *Crinum*, *Ceratophyllum*, *Blyxa*, *Elodea*, *Lagarosiphon*, and *Vallisneria*.

I used 2-gal tanks, because that is what I had on hand. However, a 5-gal tank would be much less expensive and is ideally suited for an overhead Clamp Light.

In this article, I describe two ways to keep planted tanks for pet shrimp. The bowls are easy and simple—a nice, inexpensive way for beginners to start. The DSM (Dry Start Method) is more challenging, but it is an interesting and worthy alternative to ordinary setups.

Diana Walstad is the author of *Ecology of the Planted Aquarium*. First published in 1999, the book's Fourth Edition (2023) is now available globally as a paperback and as an e-Book from Amazon. For more information on other vendors and the book, visit:

<http://dianawalstad.com/aquariums>

REFERENCES

1. Kasselmann, Christel. 2003. *Aquarium Plants*. (Krieger Publishing; Malabar, FL), p. 51.
2. Barr, Tom. 2009. Dry-Start Method. *Freshwater and Marine Aquarium* (Dec issue), pp. 54-61.
3. Barko JW and Smart RM. 1983. Effects of organic matter additions to sediment on the growth of aquatic plants. *Journal of Ecology* 71: 161-175.
4. Jablonska A *et al.* 2018. First record and DNA barcodes of the aquarium shrimp, *Neocaridina davidi*, in Central Europe from thermally polluted River Oder canal, Poland. *Knowl. Manag Aquat Ecosyst* 419, 14.
5. Vazquez ND.... LS Lopez Greco. 2017. Density and gender segregation effects in the culture of the caridean ornamental red cherry shrimp, *Neocaridina davidi* Bouvier, 1904 (Caridea: Atyidae). *J Crustacean Biol* 37: 367-373.
6. Sand-Jensen K and Borum J. 1991. Interactions among phytoplankton, epiphyton, and macrophytes in temperate freshwaters and estuaries. *Aquatic Botany* 41: 137-175.
7. Simpson PS and Eaton JW. 1986. Comparative studies of the photosynthesis of the submerged macrophyte *Elodea canadensis* and filamentous algae *Cladophora glomerata* and *Spirogyra* sp. *Aquatic Botany* 24: 1-12.